

УДК: 616.13-004.6-06:616.72-002:616.517

The role of traditional risk factors and inflammation activity in atherosclerosis development in psoriatic arthritis patients

L. V. Khimion, A. V. Boiko

Shupyk National Medical Academy of Postgraduate Education, Kyiv

It is known that patients with autoimmune diseases (rheumatoid arthritis, systemic lupus erythematosus) have early atherosclerosis development which might lead to the life threatening complications and death. Epidemiology studies demonstrated the risk of myocardial infarction, cardiovascular death is increased in patients with psoriatic arthritis (PsA), which also can be explained by premature atherosclerosis development but its risk factors (RF) remain unclear.

The objective: defining relationship of traditional RF for cardiovascular diseases (CVD) with the activity of the inflammatory process and atherosclerosis in patients with PsA.

Materials and methods. The study included 42 patients with PsA and 20 patients with skin psoriasis (PS) who did not have atherosclerotic CVD, diabetes, chronic kidney disease and other serious diseases and did not take statins. The control group consisted of 20 healthy persons who had no signs of rheumatic, infectious and other inflammatory diseases.

All enrolled patients underwent complete physical examinations, evaluation of disease activity (DAS28, PASI indexes); laboratory investigations included full blood count and ESR, CRP, plasma lipids and uric acid level. Ultrasound of carotid arteries was performed to evaluate the surrogate atherosclerosis markers – intima-media complex thickness (IMT) and atherosclerotic plaques (AP).

Results. It was determined that in patients with PsA, the severity of the atherosclerotic process is more prominent compared to patients with PS (the traditional RF profiles were the same).

Conclusion. The highest level of C-reactive protein, fibrinogen, uric acid, thicker intima-media complex was found in PsA group, which may indicate a pathogenetic association of additional RF with the development of a more aggressive atherosclerotic process.

Key words: psoriatic arthritis, psoriasis, cardiovascular diseases, dyslipidemia, atherosclerosis, C-reactive protein, uric acid.

Роль традиційних факторів ризику і активності запального процесу в розвитку атеросклерозу у хворих на псоріатичний артрит

Л.В. Хіміон, А.В. Бойко

Відомо, що для хворих на аутоімунні захворювання (ревматоїдний артрит, системний червоний вовчак) характерним є розвиток передчасного атеросклерозу, що може призводити до розвитку загрозливих для життя ускладнень та смерті. Епідеміологічні дослідження вказують на збільшений ризик інфаркту міокарда, серцево-судинної смерті серед хворих на псоріатичний артрит (ПсА), що можна пояснити передчасним атеросклерозом, однак його фактори ризику (ФР) залишаються невизначеними.

Мета дослідження: визначення взаємозв'язків між традиційними серцево-судинними факторами ризику і активністю запального процесу та атеросклерозом у хворих на ПсА.

Матеріали та методи. У дослідження було включено 42 хворих на ПсА і 20 хворих на шкірний псоріаз (ПС) без серцево-судинних захворювань, цукрового діабету, хронічної хвороби нирок, інших серйозних захворювань, які не приймали статини. До контрольної групи увійшли 20 практично здорових осіб, зіставних за статтю і віком.

Усім включеним у дослідження пацієнтам проводили повне фізикальне обстеження, визначення активності захворювання (DAS28, PASI); лабораторні дослідження (загальний аналіз крові, визначення рівню СРБ, ліпідів, сечової кислоти в сироватці крові). З метою оцінювання сурогатних маркерів атеросклерозу – товщини комплексу інтима-медіа (ТКІМ) та наявності атеросклеротичних бляшок (АБ) проводили ультразвукове обстеження сонних артерій (СА).

Результати. Встановлено більшу вираженість атеросклеротичного процесу у хворих на ПсА порівняно із хворими на ПС та групою контролю за відсутності відмінностей у поширеності традиційних факторів ризику.

Заключення. Більше значення ТКІМ СА у хворих на ПсА було пов'язано із найвищим рівнем СРБ, фібриногену, сечової кислоти, що може бути відображенням патогенетичного зв'язку додаткових факторів ризику з агресивним атеросклеротичним процесом у даного контингенту хворих.

Ключові слова: псоріатичний артрит, псоріаз, серцево-судинні захворювання, атеросклероз, дисліпідемія, СРБ, сечова кислота.

Роль традиционных факторов риска и активности воспалительного процесса в развитии атеросклероза у больных псоріатическим артритом

Л.В. Химион, А.В. Бойко

Известно, что для больных аутоиммунными заболеваниями характерным является развитие преждевременного атеросклероза, что может приводить к развитию угрожающих жизни осложнений и смерти. Эпидемиологические исследования показывают повышенный риск развития инфаркта миокарда, сердечно-сосудистой смерти среди больных псоріатическим артритом (ПсА), что можно объяснить ранним атеросклерозом, однако его факторы риска (ФР) остаются неопределенными.

Цель исследования: исследование взаимосвязей между традиционными ФР сердечно-сосудистых заболеваний и активностью воспалительного процесса у больных ПсА.

Материалы и методы. В исследование было включено 42 больных ПсА и 20 пациентов с кожным псоріазом (ПС) без сердечно-сосудистых заболеваний, сахарного диабета, хронической болезни почек, других серьезных заболеваний, не принимавших статины. В качестве группы контроля обследованы 20 практически здоровых пациентов сравнимого возраста и пола.

Всем включенным в исследование пациентам проводили полное физикальное обследование, определение активности заболевания

(DAS28, PASI); лабораторные исследования (общий анализ крови, определение уровня СРБ, липидов, мочевой кислоты в сыворотке крови). С целью определения суррогатных маркеров атеросклероза – толщины комплекса интима-медиа (ТКИМ) и наличия атеросклеротических бляшек проводили ультразвуковое исследование сонных артерий (СА).

Результаты. Выявлено большую выраженность атеросклеротического процесса у больных ПсА по сравнению с больными ПС и группой контроля при отсутствии значимых различий в профиле традиционных факторов риска.

Заключение. Большее значение ТКИМ СА у больных ПсА было связано с высоким уровнем СРБ, фибриногена, мочевой кислоты, что может быть отображением патогенетической связи дополнительных факторов риска с агрессивным атеросклеротическим процессом у данного контингента больных.

Ключевые слова: псориатический артрит, псориаз, сердечно-сосудистые заболевания, дислипидемия, атеросклероз, СРБ, мочевая кислота.

Psoriatic arthritis (PsA) is an inflammatory disease of the joints, entheses and spine, associated with psoriasis [1]. Data on the frequency of PsA vary widely and make up 0,04–3% [2, 3, 4]. Most often, the disease begins at the age of 20 to 50 years. The incidence in men and women is the same [1, 5, 12]. The prevalence of PsA among patients with psoriasis (PS) is 5,4–34% or more [6, 7, 8, 12]. In turn, the population prevalence of PS in the vast majority of studies is estimated at an average of 0,5–5% with significant fluctuations depending on the climatic-geographical region [7, 9, 10].

The study of the connection of PS and cardiovascular diseases (CVD) is paid much attention by domestic and foreign scientists. The results of clinical studies published in recent years, suggest a pathogenetic relationship between them, which is multifactorial [Ahlehoff O. et al, 2011]. In particular, it has been shown that PS and CVD are common risk factors (RF). Much attention is paid to the study of general mechanisms for the development of PS and diseases of the cardiovascular system [Balci DD et al 2009].

It has been established that up to 50% of deaths among patients with PSA are due to atherosclerosis-dependent CVD. Thus, according to the data of the meta-analysis of data [11], in 55% of patients with PsA there was an increased risk of developing CVD in this group of patients in relation to the population as a whole, whereas the overall CVD in patients with arthritis was 45% higher than the average population rate, in particular, the risk of developing myocardial infarction (MI) was higher by 68%, cerebrovascular disease – 22%, and heart failure – 31% [9, 10]. Most authors believe that this is due to the presence of atherogenic dyslipidemia (DLP) and systemic inflammation in patients with PsA.

Attention is drawn to the presence of association of cardiovascular catastrophes with an increase in serum sickness of many mediators, which are traditionally used to assess the activity of the inflammatory process, primarily C-reactive protein (CRP) [13]. Even a slight increase in CRP concentrations in a few months and years, reflecting the activity of the inflammatory process characteristic of the PS and PSA, may be a RF for subclinical inflammation in the vascular wall associated with the atherosclerotic process. In this regard, the determination of the level of CRP with the help of a highly sensitive method (the so-called hs-CRP) allows to specify the risk of development, recurrence and progression of vascular complications of atherosclerosis [13, 14].

In addition to CRP, the role of uric acid (UA) as an unconventional RF for CVD and its complications is studied at the present stage. It is stipulated that among patients with PS from 19 to 30% there is hyperuricemia, the level of which depends on the wounds of the area affected by the skin and. Titov V.N. and colleagues. (2013) have shown that in subjects with low and average risk on the SCORE scale, the concentration of uric acid (UA) in serum is associated with an average correlation with lipid profiles, regardless of the presence or absence of a metabolic syndrome, with the relationship of the UA content with triglycerides (TG) (positive) and with total cholesterol of high density lipoprotein (HDL) (negative) was detected in all studied groups [11, 15].

In the general population, the frequency of asymptomatic elevation of the UA is 5–7%, according to other data – up to

12–25%. Important clinical associations of hyperuricemia are DLP, hypertension, obesity, diabetes mellitus, gallstone disease and nephrolithiasis. Asymptomatic hyperuricemia is detected in 25% of patients with hypertension, including 75% of patients with severe hypertension (SHEP, 2000) [16]. Taking into account the «traditional» RF of CVD, obesity and metabolic syndrome, the researchers found that the level of UA 6,8–11,1 mg/dl (more than 400 $\mu\text{mol/l}$) increased 3,29 times the probability of a high level of TCh and 1,52 times – an increased amount of CRP in serum [18]. The elevated UA stimulates the activation of RAAS, enhances endothelial dysfunction and provokes hypertension. In a large meta-analysis based on 18 studies, which included 55 607 subjects with an average follow-up period of up to 6 years, it was found that the presence of asymptomatic hyperuricemia increases the risk of hypertension by 40% (RR 1,41; 95% CI: 1,23–1,58) [17].

In modern literature, there is a rather large number of studies devoted to the search for additional RFs of CVD, the definitions of which may improve the assessment and prognosis in patients with PsA with different levels of cardiovascular risk. However, until now, there is no clear answer to many of the key issues that could significantly improve the effectiveness of prevention programs and significantly reduce the high mortality and morbidity rates of CVD in patients with PsA.

The objective: determine the features of atherosclerotic process in patients with PsA.

MATERIALS AND METHODS

The study included 42 patients with PsA who did not have atherosclerotic CVD, diabetes mellitus, chronic kidney disease, and other severe illness, did not take statins. As a comparison group, we studied 20 people with PS, same age and gender and as a control group – 20 healthy persons who had no signs of rheumatic, infectious and other inflammatory diseases. The selection of the control group was done according to the gender and age of the examined patients. The selected patients, at the time of inclusion in the study, had no hypertension and did not take anti-hypertensive medications. All patients were provided with a complex of clinical and instrumental and laboratory examinations, a survey.

Anthropometry, blood pressure measurements, questionnaires for the detection of bad habits (smoking, alcohol abuse), anthropometry were used to detect FR, and in order to assess the presence of depression and / or anxiety disorders, patients used the Hospital Alert and Depression Scale (HADS), which was designed to initially detect anxiety and depression in patients (screening) in general medical practice. The burden of heredity was determined by the presence of an atherosclerotic disease or a major RF (high blood pressure, diabetes, and DLP) in relatives of the first-line patient (mother or father) who manifested before the age of 55 in men and women up to the age of 65 years. The PASI (Psoriasis Area Severity Index) calculator was used to assess the degree of skin lesions.

All patients with PsA determined the Disease Activity Score (DAS28) index for CRP, and was assessed at an average level of two years. The Alcohol Use Disorder (AUDIT) questionnaire was used for determining the risk group and the number of people

Table 1

Clinico-demographic characteristics of patients included in the study

Indicator	PsA, n=42	PS without PsA, n=20	Control group, n=20
Age, years	39,62±4,8	32,3±5,63	30,75±2,13
Women,(n/%)	20 (46,61%)	8 (40%)	11 (55%)
Men,(n/%)	22 (52,38%)	12 (60%)	9 (45%)
Average duration of the disease, years	10,42±0,34*	4,1±0,57	-
DAS 28 < 2,6	4 (8,5%)	-	-
2,6 ≤ DAS 28 < 3,2	9 (21,42%)	-	-
3,2 ≤ DAS 28 < 5,2	22 (52,38%)	-	-
DAS 28 ≥ 5,2	7 (16,66%)	-	-
PASI, points	14,36±1,12*	7,2±1,05	-
Low activity level < 10 points n, %	8 (19,04%)	3	-
Moderate degree of activity ≥ 10 < 20 points n, %	27 (64,28%)	13	-
High degree of activity ≥ 20 points n, %	7 (16,66%)	3	-

Note: * the difference between the groups is significant p < 0.05.

Table 2

Cardiovascular risk factors and state of the vascular wall in the examined patients

Indicator	PsA, n=42	PS without PsA, n=20	Control group, n=20
Smoking (at any time in the last 10 years), n/%	29/69,04*	10/52,63%#	8/40,0
Low level of physical activity, n/%	19/45,23	9/47,3%*	6/30,0
Obesity, n/%	7/16,66	3/15,78	1/5,0%
High stress lifestyle, n/%	39/92,85*	16/84,2%#	10/50,0
Family history of early CVD, n/%	29/69,09*	8/42,1	8/40,0
DLP, n/%	35/83,3*	20/62,5	3/15,0%
TC, mmol/l	5,37±0,20*	4,17±0,44	3,2±0,10
TG, mmol/l	2,45±0,22*	1,52±0,10#	1,32±0,20
HDL cholesterol, mmol/l	1,19±0,11	0,76±0,10#	1,32±0,10
LDL cholesterol, mmol/l	3,45±0,80*	2,43±0,3#	3,12±0,32
VLDL cholesterol, mmol/l	0,81±0,41**	0,5±0,13#	0,62±0,12
Index of atherogenicity, c.u.	3,69±0,91**	2,47±0,21#	2,7±0,20
CRP, mg/l	8,5±1,79**	4,45±0,53#	0,77±0,20
UA, mcmol/l	402,85±15,24**	326,68±15,59#	187,83±8,20
SCORE	3,11±0,33**	2,43±0,11#	1,71±0,11
<2 CVD RF, %	13,05%*	15,1%	14,9%
3–5 CVD RF, %	69,56%*	68,1%	67,9%
>5 CVD RF, %	17,39%*	16,8%	17,2%
Average IMT CA, mm	0,93±0,02**	0,7±0,01#	0,64±0,02
IMT ≥ 0,9 mm	45,83%**	22, 58%	5%
Atherosclerotic plaques in CA (% of patients)	6,25% (2 особи)**	0	0

Note: * the difference between the groups of patients is significant, p < 0,05; # difference with rupture control is reliable, p < 0,05.

who use the health-hazardous amount of alcohol. Identification Test, developed on the basis of the WHO Cooperative Project (1989). Tobacco use was assessed during an interview with the patient in accordance with the Order of the Ministry of

Health of Ukraine dated 03.08.2012 № 601 «On Approval and Implementation of Medical-Technological Documents on Standardization of Medical Aid in the Termination of the Use of Tobacco Products».

All patients and controls were provided with a complex of clinical and laboratory and instrumental research. A general clinical examination included: a complete physical examination, a determination of the severity of psoriasis, an articular examination. Laboratory research included determination of lipid, purine and high-sensitivity CRP indices.

In order to determine the state of the vascular wall, a duplex ultrasonic scan of carotid arteries (CA) was performed for all patients. According to the recommendations of the European Society of Cardiology, scanning of the CA was carried out in three planes – two longitudinal (front and rear) and one transverse. the intima-media complex thickness (IMT) was evaluated in the zone of maximal thickening in the orientation of the scanning plane of the longitudinal axis of the vessel. It was calculated the average value of the IMT of the right and left general CA as the mean of 9 measurements in 3 positions; the diagnostic criterion for thickening IMT was considered to be $\geq 0,9$ mm, the presence of an atherosclerotic plaque, with a local thickening of TIMC of $>1,5$ mm and more, or a thickening of more than 50% or 0.5 mm relative to other areas of IMT.

The mathematical processing of the results was carried out in the IBM SPSS 20 and Statistica 6.0 programs, according to the GCP prior to the data processing. The results base and data preparation for mathematical processing were performed in MS Excel 2007.

RESULTS

The average age of patients with PsA was $39,62 \pm 5,8$ years (20 (46,61%) women and 22 (52,38%) men), patients with PS (comparison group) $32,3 \pm 5,63$ years (8 (40%) women and 12 (60%) men respectively), the control group included 11 (55%) women and 9 (45%) men, mean age $32,3 \pm 5,63$, respectively. Duration of PSA varied from 3 to 20 years. (Table 1).

The main clinical and demographic characteristics of the examined patients are presented in Table 1.

According to the results of the initial survey, the frequency of the detection of traditional and behavioral risk factors for CVD (smoking, alcohol abuse, sedentary lifestyle) in the PSA group was not significantly different from the group of patients with skin lesions and the control group. The frequency of combination of risk factors for CVD in patients with PsA is given in Table 2.

According to the results of the analysis of lipid metabolism, a significant difference was observed between the mean levels of HF in the group of patients with PSA (Table 2) compared with the control group. The median levels of CF, TG, LDL cholesterol, LDL cholesterol were significantly higher in patients with PsA, and HDL-C levels were significantly lower than those in the FP and control groups ($p < 0,05$).

The increase in the levels of high-sensitivity CRP and SC in the group of patients with PsA was noted in 78,82% (18 persons) and 21,73% (5 persons) respectively (Table 2).

As it can be seen from the data presented in Table 2, the frequency of combining behavioral CVD risk factors was not different in the examined groups of patients (PsA and PS) compared with the control group. At the same time, in the group

of patients with PsA and PS, dyslipidemia was found to be significantly more frequent in 83,3% of PsA and 62,5% of PS and 15% of control group respectively ($p < 0,05$).

According to the results of the survey, the risk of fatal cardiovascular events by SCORE scale was determined (Table 2), the mean score of the study group was $3,11 \pm 0,33$ (moderate risk).

In the comparative analysis of lipid metabolism indexes, it was found that the average levels of TCh, TG, and LDL cholesterol in patients with PsA significantly exceeded the corresponding indices in patients with PS and indicators of healthy individuals in the control group. In this correlation analysis, the greatest influence on TIMC CA was found on DAS 28, TCh, LDL cholesterol, CRP, FG, PASI ($r = 0,54; 0,68; 0,67; 0,53; 0,52; 0,43$ in accordance). In the group of patients with DAS 28 of medium and high activity ($\geq 3,2$), a strong correlation between the IMT CA and the level of UA ($r = 0,61$) was found. The level of HDL cholesterol was associated with a strong feedback with CRP and a median power of reverse linkage with UA levels. There was a direct correlation between the level of FG and CRP and UA in the average strength ($r = 0,54; 0,48$, respectively). The given analysis in the group of patients with PS revealed: a direct connection of average strength between IMT CA, TC, LDL cholesterol, CRP ($r = 0,37; 0,39; 0,38$, respectively). The results of the correlation analysis confirmed the relationship between inflammation and dyslipidemia, which plays a significant role in the progression of atherosclerotic vascular lesions and the subsequent formation of CVD.

According to the results of ultrasound examination of carotid arteries, it was found that patients with PsA and PS have a greater severity of atherosclerotic process than healthy peers, which is confirmed by the higher average values of IMT in the group of patients with PsA ($0,93 \pm 0,02$) $p < 0,05$ compared with the group control ($0,68 \pm 0,02$), with IMT $\geq 0,9$ mm detected in 45,83% of patients with PsA, which is proven complementary RF CDV and in 22,58% of patients with PS, which is significantly higher than the control group indicator (5%). It should be noted that the presence of atherosclerotic plaques is diagnosed in 6,25% of patients with PsA.

CONCLUSIONS

1. The prevalence of behavioral risk factors for CVD and their combinations is not different in the groups of patients with PsA, PS and control group.
2. DLP is more common in patients with PsA than in patients with psoriasis ($p < 0,05$) and in the control group, which is manifested by increased levels of TC, LDL cholesterol and lower HDL cholesterol in patients with PsA compared to other subjects.
3. Patients with PsA have more aggressive atherosclerotic process comparing to patients with skin psoriasis and healthy persons, with greater average value of IMT, more frequent IMT $\geq 0,9$ mm, found at 45,83% patients and atherosclerotic plaques found in 6,25%.
4. Significant role in the development of atherosclerotic lesions in patients with PsA and PS play inflammatory factors, associated with the course of the immune-inflammatory process characteristic for psoriatic disease.

Сведения об авторах

Химион Людмила Викторовна – Национальная медицинская академия последидипломного образования имени П.Л. Шупика, 04112, г. Киев, ул. Дорогожицкая, 9

Бойко Алина Владимировна – Национальная медицинская академия последидипломного образования имени П.Л. Шупика, г. Киев, ул. Дорогожицкая, 9; тел.: (068) 388-87-76. E-mail: alina181290@gmail.com.

LITERATURE

1. Badokin V.V. Terapevicheskaya aktivnost' i bezopasnost' artrofoona pri psoriaticheskom artrite / V.V. Badokin, Yu.L. Korsakova // Consilium Medicum [E'lektronnyj resurs]. – 2006. – № 8. – Rezhim dostupa: http://con-med.ru/magazines/consilium_medicum/212569/212305/. – Data dostupa: 27.03.2014.
2. Cytokine profiles during infliximab monotherapy in psoriatic arthritis / A. Mastroianni [et al.] // British Journal of Dermatology. – 2005. – Vol. 153, № 3. – P. 531–536.
3. Gottlieb A.B. Psoriasis comorbidities / A.B. Gottlieb, C. Chao, F. Dann // Journal of Dermatological Treatment. – 2008. – Vol. 19, № 1. – P. 5–21.

4. Milosavljevic, J. Ultrasound and Power Doppler Evaluation of the Hand and Wrist in Patients with Psoriatic Arthritis / J. Milosavljevic, U. Lindqvist, A. Elvin // *Acta Radiologica*. – 2005. – Vol. 46, № 4. – P. 374–385.
5. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics / A. Gottlieb [at al.] // *Journal of the American Academy of Dermatology*. – 2008. – Vol. 58, № 5. – P. 851–864.
6. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics / A. Gottlieb [at al.] // *Journal of the American Academy of Dermatology*. – 2008. – Vol. 58, № 5. – P. 851–864.
7. Effects of alefacept on health-related quality of life in patients with psoriasis results from a randomized, placebo-controlled phase II trial / C.N. Ellis [at al.] // *American Journal of Clinical Dermatology*. – 2003. – Vol. 4, № 2. – P. 131–139.
8. Stefano I. Echographic test and cyclosporin therapy in psoriatic arthritis / I. Stefano, R. Eugenio, R. Antonio // *Journal of the European Academy of Dermatology and Venereology*. – 2000. – Vol. 14, № 3. – P. 232–233.
9. Kochergin N.G. Itogi raboty Pervoj vseimnoy konferencii po psorizmu i psoriaticheskomu artritu / N.G. Kochergin, S.N. Kochergin, L.M. Smimova // *Russkij medicinskij zhurnal [Elektronnyj resurs]*. – 2006. – Rezhim dostupa: http://www.rmj.ru/articles_4306.htm. – Data dostupa: 03.04.2014.
10. Psoriasis and diabetes: a population-based cross-sectional study / A.D. Cohen [at al.] // *Journal of the European Academy of Dermatology and Venereology*. – 2008. – Vol. 22, № 5. – P. 585–589.
11. Titov V.N. Vzaimosvyaz mochevoy kisloty s pokazatelyamy lipidnogo obmena u lits s nizkim i srednim riskom po shkale SCORE / V.N. Titov, S.A. Boytsov, S.F. Urazalina i soavt. // *Ateroskleroz i dislipidemii*. – 2013. – № 2. – S. 31–39.
12. Chandran V. Sensitivity and specificity of the caspar criteria for psoriatic arthritis in a family medicine clinic setting / V. Chandran, C.T. Schentag, D.D. Gladman // *The Journal of Rheumatology*. – 2008. – Vol. 35, № 10. – P. 2069–2070.
13. Mitchenko O.I., Lutay M.I. Dislipidemii: diagnostika. profilaktika ta likuvannya. Metodichni rekomendatsii asotsiatsii kardiologiv Ukraini. – K.: Morion. 2011.
14. Kovalenko V.M Vpliv immunnogo statusu i lipidnikh faktoriv na progresuvannya subklinichnogo aterosklerozu i rozvitok sertsevo-sudinnikh zahvoryuvan u patsientiv iz revmatoidnim artritom // V.M. Kovalenko, L.V. Khimion, G.I. Ly-senko, O.O. Garmisch / *Ukr. tevmatol. Zhurn.* – 2011.
15. Gonzalez-Juanatey C., Llorca J., Amigo-Diaz E. [et al.] High prevalence of subclinical atherosclerosis in psoriatic arthritis patients without clinically evident cardiovascular disease or classic atherosclerosis risk factors. // *Arthritis Rheum.* – 2007. – 57 (6). – P. 1074–1080.
16. Franse LV. Serum uric acid, diuretic treatment and risk of cardiovascular events in the Systolic Hypertension in the Elderly Program (SHEP) / Franse LV, Pahor M, Di Bari M et al. // *J. Hypertens.* – 2000. – Vol. 18. – P. 1149–1154.
17. Grayson PC. Hyperuricemia and incident hypertension: a systematic review and meta-analysis. / Grayson PC, Kim SY, LaValley M et al. // *Arthritis Care Res (Hoboken)*. – 2011. – Vol. 63 (1). – P. 102–110.
18. Keenan T. Relation of uric acid to serum levels of high-sensitivity C-reactive protein, triglycerides, and high-density lipoprotein cholesterol and to hepatic steatosis. / Keenan T, Blaha MJ, Nasir K et al. // *Am J Cardiol.* – 2012. – Vol. 110 (12). – P. 1787–1792.

Статья поступила в редакцию 05.11.2019